

Vegetation Types of the South West Bushland

1. Tuart Open Woodland over Jarrah – Marri – *Banksia attenuata* –Peppermint Low Woodland : found on the hill crests and upper hill slopes.



2. Tuart Marri Open Woodland over Peppermint – Jarrah – Marri – *Banksia attenuata* Low Open Forest (Peppermint dominated) : found on the major hill swales.



3. Jarrah – Marri – *Banksia attenuata* Low Woodland: found on the hill slopes (south).



4. Marri Open Woodland over Jarrah – Marri – *Banksia attenuate* – Peppermint Low Woodland: found on the hill slopes (north).



5. Jarrah – Marri Open Woodland over Jarrah – Marri – *Banksia attenuata* Low Woodland: found on the hill slopes (east).



6. Jarrah – *Banksia attenuata* Low Woodland: found on the sandplain.



7. **Marri – *Melaleuca preissiana* – *Banksia littorea* Low Woodland: found on seasonally inundated dampland.**



8. **Marri heath (previously disturbed form old Rifle Range): found on sandy flats.**



9. Marri Open Woodland over *Banksia grandis* – *B. Attenuata* Low Woodland:
found on sandy flats.



10. Cleared Areas



Note: Information and photo's are from : Flora and Fauna Survey, Onshore Environmental Consultants, September 2007.